

Wildlife Plots - Temporary Winter Grazing (Code #190)

Soil Test Rating	Potassium			
	Low K	Medium K	High K	Very High K
	Coast: 0-60 lbs/A Pied: 0-100 lbs/A	Coast: 61-150 lbs/A Pied: 101-200 lbs/A	Coast: 151-250 lbs/A Pied: 201-350 lbs/A	Coast: 250+ lbs/A Pied: 350+ lbs/A
Phosphorus	<i>Recommended Pounds N-P₂O₅-K₂O per Acre</i>			
Low P Coast: 0-30 lbs/A Pied: 0-20 lbs/A	*-100-100	*-100-50	*-100-0	*-100-0
Medium P Coast: 31-60 lbs/A Pied: 21-40 lbs/A	*-50-100	*-50-50	*-50-0	*-50-0
High P Coast: 61-100 lbs/A Pied: 41-75 lbs/A	*-0-100	*-0-50	*-0-0	*-0-0
Very High P Coast: 100+ lbs/A Pied: 75+ lbs/A	*-0-100	*-0-50	*-0-0	*-0-0

Coast = Coastal Plain Pied = Piedmont, Mountain, and Limestone Valley

Recommendations:

Recommended pH:	6.0. If the pH is less than 6.0, see Lime Table C.								
Magnesium:	If soil test Mg level is low and lime is recommended, use dolomitic limestone; if soil test Mg is low and lime is not recommended, apply 25 pounds of Mg/Acre.								
	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>Coastal Plain</td> <td>Low: 0 - 30 lbs/acre</td> <td>Medium: 31 - 60 lbs/acre</td> <td>High: >60 lbs/acre</td> </tr> <tr> <td>Piedmont</td> <td>Low: 0 - 60 lbs/acre</td> <td>Medium: 61 - 120 lbs/acre</td> <td>High: >120 lbs/acre</td> </tr> </table>	Coastal Plain	Low: 0 - 30 lbs/acre	Medium: 31 - 60 lbs/acre	High: >60 lbs/acre	Piedmont	Low: 0 - 60 lbs/acre	Medium: 61 - 120 lbs/acre	High: >120 lbs/acre
Coastal Plain	Low: 0 - 30 lbs/acre	Medium: 31 - 60 lbs/acre	High: >60 lbs/acre						
Piedmont	Low: 0 - 60 lbs/acre	Medium: 61 - 120 lbs/acre	High: >120 lbs/acre						

Fact Sheet:

Temporary winter grazing - (small grains - rye, wheat, oats). These crops can utilize about 100 pounds of nitrogen per acre during the growing season. Split the nitrogen (N) application, applying 50 pounds nitrogen per acre at planting and 50 pounds nitrogen per acre in late winter if spring grazing is desired.

Temporary winter grazing - (ryegrass alone or small grain-ryegrass). Apply 50 pounds nitrogen per acre in the fall at planting, 50 pounds per acre in late winter, and 50 pounds per acre in early spring. Ryegrass has a longer than normal grazing season. The spring and winter application of nitrogen will help extend the grazing period if that is desired.


Temporary winter grazing - (annual clovers - crimson, arrowleaf with small grains or ryegrass). Apply 30 to 50 pounds nitrogen per acre at planting. Higher rates of nitrogen will stimulate rapid grass growth and shade out the clover. If the stand contains less than 40 to 50% clover, apply 50 pounds nitrogen per acre in late winter or early spring to extend the grazing period. If only clover or other legumes such as lespedeza or vetch are planted, inoculate the seed and omit the nitrogen fertilizer.

Wildlife Plots - Temporary Winter Grazing (Code 190)

V471 - \1IX


P Recommendations, Coastal Plain

$$P_2O_5 = 127 - 1.886P + 0.00366P^2$$




K Recommendations, Coastal Plain

$$K_2O = 123 - 0.779K + 0.00083K^2$$



P Recommendations, Piedmont

$$P_2O_5 = 129 - 3.074P + 0.01435P^2$$



K Recommendations, Piedmont

$$K_2O = 128 - 0.588K + 0.00044K^2$$

